

WHAT IS CLAIMED IS:

Sub C1
1. A method of cultivating a mammalian cell in suspension *in vitro*, comprising:
(a) obtaining a mammalian cell to be cultivated in suspension;
and
(b) contacting said cell with a serum-free cell culture medium comprising at least one polyanionic or polycationic compound, wherein said medium supports the cultivation of said cell in suspension.

10 2. The method of claim 1, wherein said polyanionic compound is a polysulfonated compound or a polysulfated compound.

Sub E2
3. The method of claim 2, wherein said polysulfonated or polysulfated compound is selected from the group consisting of dextran sulfate, heparin, heparan sulfate, chondroitin sulfate, dermatan sulfate, pentosan sulfate and a proteoglycan.

15 4. The method of claim 2, wherein said polysulfonated or polysulfated compound is dextran sulfate.

5. The method of claim 4, wherein said dextran sulfate has an average molecular weight of about 5,000 daltons.

6. The method of claim 4, wherein said medium is protein-free.

20 7. The method of claim 1, wherein said medium is a 1X medium formulation.

8. The method of claim 1, wherein said medium formulation is a 10X concentrated medium formulation.

5
Sub F17
9. The method of claim 1, wherein said medium further comprises one or more ingredients selected from the group of ingredients consisting of one or more amino acids, one or more vitamins, one or more inorganic salts, one or more buffering salts, one or more sugars, one or more lipids, transferrin (or transferrin substitute), and insulin (or insulin substitute).

10
10. The method of claim 9, wherein said medium further comprises one or more supplements selected from the group of supplements consisting of one or more cytokines, heparin, one or more animal peptides, one or more yeast peptides and one or more plant peptides.

11. The method of claim 10, wherein said one or more plant peptides are one or more rice peptides or one or more soy peptides.

15
12. The method of claim 9, wherein said amino acid ingredient comprises one or more amino acids selected from the group consisting of L-alanine, L-arginine, L-asparagine, L-aspartic acid, L-cysteine, L-glutamic acid, L-glutamine, glycine, L-histidine, L-isoleucine, L-leucine, L-lysine, L-methionine, L-phenylalanine, L-proline, L-serine, L-threonine, L-tryptophan, L-tyrosine and L-valine.

20
13. The method of claim 9, wherein said vitamin ingredient comprises one or more vitamins selected from the group consisting of biotin, choline chloride, D-Ca⁺⁺-pantothenate, folic acid, *i*-inositol, niacinamide, pyridoxine, riboflavin, thiamine and vitamin B₁₂.

14. The method of claim 9, wherein said inorganic salt ingredient comprises one or more inorganic salts selected from the group consisting of one or more calcium salts, $\text{Fe}(\text{NO}_3)_3$, KCl , one or more magnesium salts, one or more manganese salts, NaCl , NaHCO_3 , Na_2HPO_4 , one or more selenium salts, one or more vanadium salts and one or more zinc salts.

15. A method of cultivating a mammalian cell in suspension *in vitro*, comprising:

(a) obtaining a mammalian cell to be cultivated in suspension; and

(b) contacting said cell with a cell culture medium comprising the ingredients ethanolamine, D-glucose, N-[2-hydroxyethyl]piperazine-N'-[2-ethanesulfonic acid] (HEPES), insulin, linoleic acid, lipoic acid, phenol red, PLURONIC F68, putrescine, sodium pyruvate, transferrin, L-alanine, L-arginine, L-asparagine, L-aspartic acid, L-cysteine, L-glutamic acid, L-glutamine, glycine, L-histidine, L-isoleucine, L-leucine, L-lysine, L-methionine, L-phenylalanine, L-proline, L-serine, L-threonine, L-tryptophan, L-tyrosine, L-valine, biotin, choline chloride, D- Ca^{++} -pantothenate, folic acid, *D*-inositol, niacinamide, pyridoxine, riboflavin, thiamine, vitamin B_{12} , at least one polyanionic or polycationic compound, one or more calcium salts, KCl , one or more iron salts, one or more magnesium salts, one or more manganese salts, NaCl , NaHCO_3 , Na_2HPO_4 , one or more selenium salts, one or more vanadium salts and one or more zinc salts, wherein each ingredient is present in an amount which supports the cultivation of said cell in suspension.

16. The method of claim 15, wherein said polyanionic compound is a polysulfonated or polysulfated compound.

Sub E4
17. The method of claim 16, wherein said polysulfonated or polysulfated compound is selected from the group consisting of dextran sulfate, heparin, heparan sulfate, chondroitin sulfate, dermatan sulfate, pentosan sulfate and a proteoglycan.

5 18. The method of claim 17, wherein said polysulfonated or polysulfated compound is dextran sulfate.

19. The method of claim 18, wherein said dextran sulfate has an average molecular weight of about 5,000 daltons.

10 20. The method of claim 15, wherein said medium further comprises one or more supplements selected from the group consisting of one or more cytokines, heparin, one or more animal peptides, one or more yeast peptides and one or more plant peptides.

21. The method of claim 20, wherein said one or more plant peptides are one or more rice peptides or one or more soy peptides.

15 22. A method of cultivating a mammalian cell in suspension *in vitro*, comprising:

Sub C3
(a) obtaining a mammalian cell to be cultivated in suspension;

and

20 (b) contacting said cell with a cell culture medium obtained by combining a basal medium with at least one polyanionic or polycationic compound, wherein said medium supports the cultivation of said cell in suspension.

23. The method of claim 22, wherein said polyanionic compound is a polysulfonated or polysulfated compound.

Sub E6
5 24. The method of claim 23, wherein said polysulfonated or polysulfated compound is selected from the group consisting of dextran sulfate, heparin, heparan sulfate, chondroitin sulfate, dermatan sulfate, pentosan sulfate and a proteoglycan.

25. The method of claim 23, wherein said polysulfonated or polysulfated compound is dextran sulfate.

10 26. The method of claim 25, wherein said dextran sulfate has an average molecular weight of about 5,000 daltons.

15 27. The method of claim 22, wherein said basal medium is obtained by combining one or more ingredients selected from the group consisting of ethanolamine, D-glucose, N-[2-hydroxyethyl]-piperazine-N'-[2-ethanesulfonic acid] (HEPES), insulin, linoleic acid, lipoic acid, phenol red, PLURONIC F68, putrescine, sodium pyruvate, transferrin, L-alanine, L-arginine, L-asparagine, L-aspartic acid, L-cysteine, L-glutamic acid, L-glutamine, glycine, L-histidine, L-isoleucine, L-leucine, L-lysine, L-methionine, L-phenylalanine, L-proline, L-serine, L-threonine, L-tryptophan, L-tyrosine, L-valine, biotin, choline chloride, D-Ca⁺⁺-pantothenate, folic acid, *D*-inositol, niacinamide, pyridoxine, riboflavin,
20 thiamine, vitamin B₁₂, one or more calcium salts, one or more iron salts, KCl, one or more magnesium salts, one or more manganese salts, NaCl, NaHCO₃, Na₂HPO₄, one or more selenium salts, one or more vanadium salts and one or more zinc salts,

25 wherein each ingredient is added in an amount which supports the cultivation of said cell in suspension.

28. The method of claim 22, wherein said medium is obtained by combining said basal medium and one or more supplements selected from the group consisting of one or more cytokines, heparin, one or more animal peptides, one or more yeast peptides and one or more plant peptides.

5 29. The method of claim 28, wherein said one or more plant peptides are one or more rice peptides or one or more soy peptides.

30. The method of any one of claims 1, 15 or 22, wherein said mammalian cell is a mammalian epithelial cell.

10 31. The method of claim 30, wherein said mammalian epithelial cell is selected from the group consisting of a keratinocyte, a cervical epithelial cell, a bronchial epithelial cell, a tracheal epithelial cell, a kidney epithelial cell and a retinal epithelial cell.

32. The method of claim 30, wherein said cell is a human cell.

15 33. The method of claim 32, wherein said human cell is a 293 embryonic kidney cell, a HeLa cervical epithelial cell, a PER-C6 retinal cell, or a derivative thereof.

34. The method of claim 33, wherein said human cell is a 293 embryonic kidney cell.

35. The method of claim 30, wherein said cell is a normal cell.

20 36. The method of claim 30, wherein said cell is an abnormal cell.

00023514-022398
362220-47532060

37. The method of claim 36, wherein said abnormal cell is a transformed cell, an established cell, or a cell derived from a diseased tissue sample.

38. A kit for the cultivation of a mammalian epithelial cell in suspension *in vitro*, said kit comprising one or more containers, wherein a first container contains a serum-free culture medium comprising at least one polyanionic or polycationic compound, wherein said medium supports the cultivation of said cell in suspension.

39. A kit for the cultivation of a mammalian epithelial cell in suspension *in vitro*, said kit comprising one or more containers, wherein a first container contains a cell culture medium comprising the ingredients ethanolamine, D-glucose, N-[2-hydroxyethyl]piperazine-N'-[2-ethanesulfonic acid] (HEPES), insulin, linoleic acid, lipoic acid, phenol red, PLURONIC F68, putrescine, sodium pyruvate, transferrin, L-alanine, L-arginine, L-asparagine, L-aspartic acid, L-cysteine, L-glutamic acid, L-glutamine, glycine, L-histidine, L-isoleucine, L-leucine, L-lysine, L-methionine, L-phenylalanine, L-proline, L-serine, L-threonine, L-tryptophan, L-tyrosine, L-valine, biotin, choline chloride, D-Ca⁺⁺-pantothenate, folic acid, *i*-inositol, niacinamide, pyridoxine, riboflavin, thiamine, vitamin B₁₂, one or more calcium salts, KCl, one or more iron salts, one or more magnesium salts, one or more manganese salts, NaCl, NaHCO₃, Na₂HPO₄, one or more selenium salts, one or more vanadium salts and one or more zinc salts,

wherein each ingredient is present in an amount which supports the cultivation of said cell in suspension.

40. The kit of claim 39, wherein said culture medium in said first container further comprises at least one polyanionic or polycationic compound.

00028514.022396
962220.47582060

41. The kit of claim 39, further comprising a second container containing at least one polyanionic or polycationic compound.

42. The kit of any one of claims 38, 40 or 41, wherein said polyanionic compound is a polysulfonated or polysulfated compound.

43. The kit of claim 42, wherein said polysulfonated or polysulfated compound is selected from the group consisting of dextran sulfate, heparin, heparan sulfate, chondroitin sulfate, dermatan sulfate, pentosan sulfate and a proteoglycan.

44. The kit of claim 42, wherein said polysulfonated or polysulfated compound is dextran sulfate.

45. The kit of claim 44, wherein said dextran sulfate has an average molecular weight of about 5,000 daltons.

46. The kit of claim 38 or claim 39, further comprising one or more additional containers containing one or more supplements selected from the group consisting of one or more cytokines, heparin, one or more animal peptides, one or more yeast peptides and one or more plant peptides.

47. The kit of claim 46, wherein said one or more plant peptides are one or more rice peptides or one or more soy peptides.

48. A composition comprising a serum-free culture medium comprising at least one polyanionic or polycationic compound, wherein said culture medium supports the cultivation of a mammalian cell in suspension *in vitro*.

09028514-022398
862220-41582060

49. The composition of claim 48, wherein said polyanionic compound is a polysulfonated or polysulfated compound.

50. The composition of claim 49, wherein said polysulfonated or polysulfated compound is selected from the group consisting of dextran sulfate, heparin, heparan sulfate, chondroitin sulfate, dermatan sulfate, pentosan sulfate and a proteoglycan.

51. The composition of claim 49, wherein said polysulfonated or polysulfated compound is dextran sulfate.

52. The composition of claim 51, wherein said dextran sulfate has an average molecular weight of about 5,000 daltons.

53. The composition of claim 48, further comprising at least one virus.

54. The composition of claim 53, wherein said virus is selected from the group consisting of an adenovirus, an adeno-associated virus and a retrovirus.

55. The composition of claim 48, further comprising at least one mammalian epithelial cell.

56. The composition of claim 55, wherein said mammalian epithelial cell is selected from the group consisting of a keratinocyte, a cervical epithelial cell, a bronchial epithelial cell, a tracheal epithelial cell, a kidney epithelial cell and a retinal epithelial cell.

57. The composition of claim 55, wherein said cell is a human cell.

09028514-022398

5

60. The composition of claim 55, wherein said cell is an abnormal cell.

10

15

20

65. The composition of claim 63, wherein said polysulfonated or polysulfated compound is dextran sulfate.

claim 65,
5,000 dal

4

~~culture medium~~

~~culture medium~~

~~culture medium~~
~~and is a polysulfide~~

ate, chondroitin

culture medium containing
und is dextran su

15

73. A method of producing a virus comprising

(a) obtaining a mammalian cell to be infected with a virus;

(b) contacting said cell with a virus under conditions suitable to promote the infection of said cell by said virus; and

(c) cultivating said cell according to the method of any one of claims 1, 15 or 22, under conditions suitable to promote the production of said virus by said cell.

74. The method of claim 73, wherein said mammalian cell is an epithelial cell.

75. The method of claim 73, wherein said mammalian cell is a human cell.

76. The method of claim 75, wherein said human cell is a 293 embryonic kidney cell.

77. The method of claim 73, wherein said virus is an adenovirus, an adeno-associated virus or a retrovirus.

~~78. A virus produced according to the method of claim 73.~~

79. A method of producing a polypeptide comprising

(a) obtaining a mammalian cell that has been genetically engineered to produce a polypeptide; and

(b) cultivating said cell according to the method of any one of claims 1, 15 or 22, under conditions favoring expression of said polypeptide by said mammalian cell.

5

10

15

84. A eukaryotic cell culture medium comprising a Fe^{2+} chelate and a Zn^{2+} salt, wherein said medium is capable of supporting the high-density growth of mammalian cells in suspension culture and/or the expression of recombinant protein.

86. The eukaryotic cell culture medium according to claim 84, wherein said mammalian cells are Chinese hamster ovary cells.

87. The eukaryotic cell culture medium according to claim 84, wherein said medium is a 1X medium formulation.

88. The eukaryotic cell culture medium according to claim 84, wherein said medium is a concentrated medium formulation.

89. The eukaryotic cell culture medium according to claim 88, wherein said medium is a 10X medium formulation.

90. The eukaryotic cell culture medium according to claim 88, wherein said medium formulation is greater than 10X.

91. The eukaryotic cell culture medium according to claim 84, wherein the concentration of said Fe^{2+} is about 0.00028 to 0.011 g/L and said concentration of said Zn^{2+} is about 0.00007 to 0.00073 g/L.

92. The eukaryotic cell culture medium according to claim 91 wherein said concentration of said Fe^{2+} is about 0.0011 g/L and said concentration of said Zn^{2+} is about 0.000354 g/L.

93. The eukaryotic cell culture medium according to claim 84, further comprising a polyanionic or polycationic compound, wherein said polyanionic or polycationic compound is present in an amount sufficient to prevent cell clumping and/or increase the level of recombinant protein expression.

94. The eukaryotic cell culture medium according to claim 93, wherein said polyanionic compound is a polysulfonated compound or a polysulfated compound.

95. The eukaryotic cell culture medium according to claim 94, wherein said polysulfonated or polysulfated compound is selected from the group consisting of dextran sulfate, heparin, heparan sulfate, chondroitin sulfate, dermatan sulfate, pentosan sulfate and a proteoglycan.

1994-1995

51

19

2

2

5 99. A eukaryotic cell culture medium obtained by combining ferrous sulfate·EDTA and $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$, together with a eukaryotic medium, wherein said ferrous sulfate·EDTA and said $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ are each present in an amount which supports the high-density growth of Chinese hamster ovary cells in suspension culture.

100. The eukaryotic cell culture medium according to claim 99, further comprising a polyanionic or polycationic compound, wherein said polyanionic or polycationic compound is present in an amount sufficient to prevent cell clumping and/or increase the level of recombinant protein expression.

101. The eukaryotic cell culture medium according to claim 100, wherein said polyanionic compound is a polysulfonated compound or a polysulfated compound.

102. The eukaryotic cell culture medium according to claim 101, wherein said polysulfonated or polysulfated compound is selected from the group consisting of dextran sulfate, heparin, heparan sulfate, chondroitin sulfate, dermatan sulfate, pentosan sulfate and a proteoglycan.

103. The eukaryotic cell culture medium according to claim 102, wherein said polysulfonated or polysulfated compound is dextran sulfate.

104. The eukaryotic cell culture medium according to claim 103, wherein said dextran sulfate has an average molecular weight of 5,000 daltons.

105. The eukaryotic cell culture medium obtained according to claim 99 or 100, further comprising one or more ingredients selected from the group consisting of L-arginine, L-asparagine· H_2O , L-aspartic acid, L-glutamic acid, L-

histidine, hydroxy-L-proline, L-isoleucine, L-leucine, L-lysine·HCl, L-methionine, L-phenylalanine, L-proline, L-serine, L-threonine, L-tryptophan, L-tyrosine, L-valine, L-cystine·2HCl, Na₂HPO₄, pyridoxine·HCl, thiamine·HCl, glutathione, cupric sulfate·5H₂O, cadmium chloride·5H₂O, cobalt chloride·2H₂O, stannous chloride·2H₂O, manganous sulfate·H₂O, nickel sulfate·6H₂O, sodium metavanadate, ammonium molybdate·4H₂O, barium acetate, potassium bromide, potassium iodide, chromium sulfate, sodium fluoride, silver nitrate, rubidium chloride, zirconyl chloride, aluminum chloride, germanium dioxide, titanium tetrachloride, sodium metasilicate, magnesium chloride (anhydrous), D-calcium pantothenate, calcium nitrate·4H₂O, potassium chloride, ascorbic acid magnesium salt phosphate, pluronic F68 10% solution, Na₂HPO₄, D-glucose, folic acid, riboflavin, biotin, choline chloride, niacinamide, i-inositol, sodium pyruvate, vitamin B-12, β-mercaptoethanol, para-aminobenzoic acid, β-glycerophosphate, sodium selenite, ethanolamine·HCl, spermine, putrescine·2HCl, monothioglycerol, and sodium bicarbonate,

wherein each of said ingredients is present in an amount which supports the high-density growth of Chinese hamster ovary cells in suspension culture and/or the expression of recombinant protein.

106. A method of cultivating mammalian cells in suspension culture to high density and/or expressing a recombinant protein, said method comprising the steps of

(a) contacting said cells with the eukaryotic cell culture medium of claim 84, wherein said Fe²⁺ chelate and said Zn²⁺ salt are each present in an amount which supports the growth of mammalian cells in culture; and

(b) cultivating said mammalian cells under conditions suitable to support the growth of said cells to high density and/or the expression of said recombinant protein.

09025514-022398
866220-41582060

Sub-B1C4

Sub E8

107. The method according to claim 106, further comprising a polyanionic or polycationic compound, wherein said polyanionic or polycationic compound is present in an amount sufficient to prevent cell clumping and/or increase the level of recombinant protein expression.

5

108. The method according to claim 107, wherein said polyanionic compound is a polysulfonated compound or a polysulfated compound.

Sub E9

109. The method according to claim 108, wherein said polysulfonated or polysulfated compound is selected from the group consisting of dextran sulfate, heparin, heparan sulfate, chondroitin sulfate, dermatan sulfate, pentosan sulfate and a proteoglycan.

10

110. The method according to claim 109, wherein said polysulfonated or polysulfated compound is dextran sulfate.

111. The method according to claim 110, wherein said dextran sulfate has an average molecular weight of 5,000 daltons.

15

112. The method according to claim 106, or 107, wherein said eukaryotic cell culture medium further comprises one or more ingredients selected from the group consisting of L-arginine, L-asparagine·H₂O, L-aspartic acid, L-glutamic acid, L-histidine, hydroxy-L-proline, L-isoleucine, L-leucine, L-lysine·HCl, L-methionine, L-phenylalanine, L-proline, L-serine, L-threonine, L-tryptophan, L-tyrosine, L-valine, L-cystine·2HCl, Na₂HPO₄, pyridoxine·HCl, thiamine·HCl, glutathione, cupric sulfate·5H₂O, cadmium chloride·5H₂O, cobalt chloride·2H₂O, stannous chloride·2H₂O, manganous sulfate·H₂O, nickel sulfate·6H₂O, sodium metavanadate, ammonium molybdate·4H₂O, barium acetate, potassium bromide, potassium iodide, chromium sulfate, sodium fluoride,

20

0902514-022398

silver nitrate, rubidium chloride, zirconyl chloride, aluminum chloride, germanium dioxide, titanium tetrachloride, sodium metasilicate, magnesium chloride (anhydrous), D-calcium pantothenate, calcium nitrate-4H₂O, potassium chloride, ascorbic acid magnesium salt phosphate, pluronic F68 10% solution, Na₂HPO₄, D-glucose, folic acid, riboflavin, biotin, choline chloride, niacinamide, i-inositol, sodium pyruvate, vitamin B-12, β -mercaptoethanol, para-aminobenzoic acid, β -glycerophosphate, sodium selenite, ethanolamine-HCl, spermine, putrescine-2HCl, monothioglycerol, and sodium bicarbonate,

wherein each of said ingredients is present in an amount which supports the ~~high-density~~ growth of Chinese hamster ovary cells in suspension culture and/or the expression of recombinant protein.

113. An improved eukaryotic cell culture medium for growing Chinese hamster ovary cells in suspension culture to high density, the improvement comprising replacing insulin with a Zn²⁺ salt and/or replacing transferrin with a Fe²⁺ chelate and/or replacing transferrin with a Fe³⁺ chelate.

114. The eukaryotic cell culture medium according to claim 113, further comprising a polyanionic or polycationic compound, wherein said polyanionic or polycationic compound is present in an amount sufficient to prevent cell clumping and/or increase the level of recombinant protein expression.

115. The eukaryotic cell culture medium according to claim 114, wherein said polyanionic compound is a polysulfonated compound or a polysulfated compound.

116. The eukaryotic cell culture medium according to claim 115, wherein said polysulfonated or polysulfated compound is selected from the group

09028514-022998

5

10

15

20

consisting of dextran sulfate, heparin, heparan sulfate, chondroitin sulfate, dermatan sulfate, pentosan sulfate and a proteoglycan.

117. The eukaryotic cell culture medium according to claim 116, wherein said polysulfonated or polysulfated compound is dextran sulfate.

118. The eukaryotic cell culture medium according to claim 117, wherein said dextran sulfate has an average molecular weight of 5,000 daltons.

119. The eukaryotic cell culture medium according to claim 113, wherein the concentration of said Fe^{2+} or Fe^{3+} , if present, is about 0.00028 to 0.011 g/L and said concentration of said Zn^{2+} , if present, is about 0.00007 to 0.00073 g/L.

120. The eukaryotic cell culture medium according to claim 119, wherein said concentration of said Fe^{2+} or Fe^{3+} , if present, is about 0.0011 g/L and said concentration of said Zn^{2+} , if present, is about 0.000354 g/L.

121. The eukaryotic cell culture medium according to claim 113 or 114, further comprising one or more ingredients selected from the group consisting of L-arginine, L-asparagine· H_2O , L-aspartic acid, L-glutamic acid, L-histidine, hydroxy-L-proline, L-isoleucine, L-leucine, L-lysine·HCl, L-methionine, L-phenylalanine, L-proline, L-serine, L-threonine, L-tryptophan, L-tyrosine, L-valine, L-cystine·2HCl, Na_2HPO_4 , pyridoxine·HCl, thiamine·HCl, glutathione, cupric sulfate· $5\text{H}_2\text{O}$, cadmium chloride· $5\text{H}_2\text{O}$, cobalt chloride· $2\text{H}_2\text{O}$, stannous chloride· $2\text{H}_2\text{O}$, manganous sulfate· H_2O , nickel sulfate· $6\text{H}_2\text{O}$, sodium metavanadate, ammonium molybdate· $4\text{H}_2\text{O}$, barium acetate, potassium bromide, potassium iodide, chromium sulfate, sodium fluoride, silver nitrate, rubidium chloride, zirconyl chloride, aluminum chloride, germanium dioxide, titanium

5 tetrachloride, sodium metasilicate, magnesium chloride (anhydrous), D-calcium
pantothenate, calcium nitrate·4H₂O, potassium chloride, ascorbic acid magnesium
salt phosphate, pluronic F68 10% solution, Na₂HPO₄, D-glucose, folic acid,
riboflavin, biotin, choline chloride, niacinamide, i-inositol, sodium pyruvate,
vitamin B-12, β-mercaptoethanol, para-aminobenzoic acid, β-glycerophosphate,
sodium selenite, ethanolamine·HCl, spermine, putrescine·2HCl,
monothioglycerol, and sodium bicarbonate,

wherein each of said ingredients is present in an amount which supports
the high-density growth of Chinese hamster ovary cells in suspension culture
and/or the expression of recombinant protein.

122. An improved method of cultivating mammalian cells in
suspension culture to high density, said improvement comprising the steps of

(a) contacting said cells with the eukaryotic cell culture medium of
claim 84, wherein said Fe²⁺ chelate and said Zn²⁺ salt are each present in an
amount which supports the high-density growth of mammalian cells in culture;
and

(b) cultivating said mammalian cells under conditions suitable to
support the high-density growth of said cells.

123. The method according to claim 122, further comprising a
polyanionic or polycationic compound, wherein said polyanionic or polycationic
compound is present in an amount sufficient to prevent cell clumping and/or
increase the level of recombinant protein expression.

124. The method according to claim 123, wherein said polyanionic
compound is a polysulfonated compound or a polysulfated compound.

05023514-02298

125. The method according to claim 124, wherein said polysulfonated or polysulfated compound is selected from the group consisting of dextran sulfate, heparin, heparan sulfate, chondroitin sulfate, dermatan sulfate, pentosan sulfate and a proteoglycan.

126. The method according to claim 125, wherein said polysulfonated or polysulfated compound is dextran sulfate.

127. The method according to claim 126, wherein said dextran sulfate has an average molecular weight of 5,000 daltons.

128. The method according to claim 122 or 123, wherein said eukaryotic cell culture medium further comprises one or more ingredients selected from the group consisting of L-arginine, L-asparagine·H₂O, L-aspartic acid, L-glutamic acid, L-histidine, hydroxy-L-proline, L-isoleucine, L-leucine, L-lysine·HCl, L-methionine, L-phenylalanine, L-proline, L-serine, L-threonine, L-tryptophan, L-tyrosine, L-valine, L-cystine·2HCl, Na₂HPO₄, pyridoxine·HCl, thiamine·HCl, glutathione, cupric sulfate·5H₂O, cadmium chloride·5H₂O, cobalt chloride·2H₂O, stannous chloride·2H₂O, manganous sulfate·H₂O, nickel sulfate·6H₂O, sodium metavanadate, ammonium molybdate·4H₂O, barium acetate, potassium bromide, potassium iodide, chromium sulfate, sodium fluoride, silver nitrate, rubidium chloride, zirconyl chloride, aluminum chloride, germanium dioxide, titanium tetrachloride, sodium metasilicate, magnesium chloride (anhydrous), D-calcium pantothenate, calcium nitrate·4H₂O, potassium chloride, ascorbic acid magnesium salt phosphate, pluronic F68 10% solution, Na₂HPO₄, D-glucose, folic acid, riboflavin, biotin, choline chloride, niacinamide, i-inositol, sodium pyruvate, vitamin B-12, β-mercaptoethanol, para-aminobenzoic acid, β-glycerophosphate, sodium selenite, ethanolamine·HCl, spermine, putrescine·2HCl, monothioglycerol, and sodium bicarbonate,

wherein each of said ingredients is present in an amount which supports the high-density growth of Chinese hamster ovary cells in suspension culture.

129. A kit for the cultivation of a mammalian epithelial cell in suspension *in vitro*, said kit comprising one or more containers, wherein a first container contains a medium comprising a Fe^{2+} chelate and a Zn^{2+} salt, wherein said medium is capable of supporting the high-density growth of mammalian cells in suspension culture and/or the expression of recombinant protein.

130. The kit according to claim 129, wherein said medium contains neither transferrin nor insulin.

131. The kit according to claim 130, wherein said mammalian cells are Chinese hamster ovary cells.

132. The kit according to claim 129, said first container further comprising one or more ingredients selected from the group consisting of L-arginine, L-asparagine· H_2O , L-aspartic acid, L-glutamic acid, L-histidine, hydroxy-L-proline, L-isoleucine, L-leucine, L-lysine· HCl , L-methionine, L-phenylalanine, L-proline, L-serine, L-threonine, L-tryptophan, L-tyrosine, L-valine, L-cystine· 2HCl , Na_2HPO_4 , pyridoxine· HCl , thiamine· HCl , glutathione, cupric sulfate· $7\text{H}_2\text{O}$, cadmium chloride· $5\text{H}_2\text{O}$, cobalt chloride· $2\text{H}_2\text{O}$, stannous chloride· $2\text{H}_2\text{O}$, manganous sulfate· H_2O , nickel sulfate· $6\text{H}_2\text{O}$, sodium metavanadate, ammonium molybdate· $4\text{H}_2\text{O}$, barium acetate, potassium bromide, potassium iodide, chromium sulfate, sodium fluoride, silver nitrate, rubidium chloride, zirconyl chloride, aluminum chloride, germanium dioxide, titanium tetrachloride, sodium metasilicate, magnesium chloride (anhydrous), D-calcium pantothenate, calcium nitrate· $4\text{H}_2\text{O}$, potassium chloride, ascorbic acid magnesium salt phosphate, pluronic F68 10% solution, Na_2HPO_4 , D-glucose, folic acid,

riboflavin, biotin, choline chloride, niacinamide, i-inositol, sodium pyruvate, vitamin B-12, β -mercaptoethanol, para-aminobenzoic acid, β -glycerophosphate, sodium selenite, ethanolamine·HCl, spermine, putrescine·2HCl, monothioglycerol, and sodium bicarbonate,

5 wherein each of said ingredients is present in an amount which supports the high-density growth of Chinese hamster ovary cells in suspension culture and/or the expression of recombinant protein.

133. The kit according to claim 132, wherein said medium in said first container further comprises at least one polyanionic or polycationic compound.

10 134. The kit according to claim 132, further comprising a second container containing at least one polyanionic or polycationic compound.

135. The kit according to any one of claims 132, 133, or 134, wherein said polyanionic compound is a polysulfonated or polysulfated compound.

15 136. The kit according to claim 135, wherein said polysulfonated or polysulfated compound is selected from the group consisting of dextran sulfate, heparin, heparan sulfate, chondroitin sulfate, dermatan sulfate, pentosan sulfate and a proteoglycan.

137. The kit according to claim 136, wherein said polysulfonated or polysulfated compound is dextran sulfate.

20 138. The kit according to claim 137, wherein said dextran sulfate has an average molecular weight of about 5,000 daltons.

00028544.022398
86E220.4T582060

139. The method of claim 9, wherein said medium further comprises linoleic acid, lipoic acid, phenol red, PLURONIC F68, putrescine, sodium pyruvate, wherein said buffering salt is N-[2-hydroxyethyl]-piperazine-N'-[2-ethanesulfonic acid] (HEPES), and wherein said sugar is D-glucose.

Add D³

Add B²
Add C⁷

09028544-023398